

Two-loop test of the $\mathcal{N} = 6$ Chern-Simons theory S -matrix

Changrim Ahn ¹ and Rafael I. Nepomechie ²

Abstract

Starting from the integrable two-loop spin-chain Hamiltonian describing the anomalous dimensions of scalar operators in the planar $\mathcal{N} = 6$ superconformal Chern-Simons theory of ABJM, we perform a direct coordinate Bethe ansatz computation of the corresponding two-loop S -matrix. The result matches with the weak-coupling limit of the scalar sector of the all-loop S -matrix which we have recently proposed. In particular, we confirm that the scattering of $\mathcal{A}$ and $\mathcal{B}$ particles is reflectionless. As a warm up, we first review the analogous computation of the one-loop S -matrix from the one-loop dilatation operator for the scalar sector of planar $\mathcal{N} = 4$ superconformal Yang-Mills theory, and compare the result with the all-loop $SU(2|2)^2$ S -matrix.

¹ Department of Physics, Ewha Womans University, Seoul 120-750, South Korea

² Physics Department, P.O. Box 248046, University of Miami, Coral Gables, FL 33124 USA

1 Introduction

Exact factorized S -matrices [1] play a key role in the understanding of integrable models. Planar four-dimensional $\mathcal{N} = 4$ superconformal Yang-Mills (YM) theory (and therefore, according to the AdS_5/CFT_4 correspondence [2], a certain type IIB superstring theory on $AdS_5 \times S^5$) is believed to be integrable (see [3]-[6] and references therein). A corresponding exact factorized S -matrix with $SU(2|2)^2$ symmetry has been proposed (see [7]-[14] and references therein), which leads [8, 15, 16] to the all-loop Bethe ansatz equations (BAEs) [17].

Aharony, Bergman, Jafferis and Maldacena (ABJM) [18] recently proposed an analogous AdS_4/CFT_3 correspondence relating planar three-dimensional $\mathcal{N} = 6$ superconformal Chern-Simons (CS) theory to type IIA superstring theory on $AdS_4 \times CP^3$. Minahan and Zarembo [19] subsequently found that the scalar sector of $\mathcal{N} = 6$ CS is integrable at the leading two-loop order, and proposed two-loop BAEs for the full theory (see also [20]). Moreover, evidence for classical integrability of the dual string sigma model (large-coupling limit) was discovered in [21, 22, 23]. On the basis of these results, and assuming integrability to all orders, Gromov and Vieira then conjectured all-loop BAEs [24].

Based on the symmetries and the spectrum of elementary excitations [19, 25, 26], we proposed an exact factorized AdS_4/CFT_3 S -matrix [28]. As a check, we verified that this S -matrix leads to the all-loop BAEs in [24]. An unusual feature of this S -matrix is that the scattering of $\mathcal{A}$ and $\mathcal{B}$ particles is reflectionless. (A similar S -matrix which is not reflectionless is not consistent with the known two-loop BAEs [29].) For further related developments of the AdS_4/CFT_3 correspondence, see [30] and references therein.

Considerable guesswork has entered into the above-mentioned all-loop results. While there is substantial evidence for the all-loop BAEs and S -matrix in the well-studied AdS_5/CFT_4 case, the same cannot be said for the rapidly-evolving AdS_4/CFT_3 case.

In an effort to further check our proposed S -matrix, we perform here a direct coordinate Bethe ansatz computation of the two-loop S -matrix, starting from the integrable two-loop spin-chain Hamiltonian describing the anomalous dimensions of scalar operators in planar $\mathcal{N} = 6$ CS [19]. The result matches with the weak-coupling limit of the scalar sector of our all-loop S -matrix [28]. In particular, we confirm that the scattering of $\mathcal{A}$ and $\mathcal{B}$ particles is reflectionless. As a warm up, we first review the analogous computation by Berenstein and Vázquez [5] of the one-loop S -matrix from the one-loop dilatation operator for the scalar sector of planar $\mathcal{N} = 4$ YM [3], and compare the result with the all-loop $SU(2|2)^2$ S -matrix.

The outline of this paper is as follows. In Sec. 2 we review the simpler case of $\mathcal{N} = 4$ YM. In Sec. 3 we analyze the $\mathcal{N} = 6$ CS case, relegating most of the details of $\mathcal{A} - \mathcal{B}$ scattering

to an appendix. We briefly discuss our results in Sec. 4.

2 One-loop S -matrix in the scalar sector of $\mathcal{N} = 4$ YM

As is well known, $\mathcal{N} = 4$ YM has six scalar fields $\Phi_i(x)$ ($i = 1, \dots, 6$) in the adjoint representation of $SU(N)$. It is convenient to associate single-trace gauge-invariant scalar operators with states of an $SO(6)$ quantum spin chain with L sites,

$$\text{tr } \Phi_{i_1}(x) \cdots \Phi_{i_L}(x) \quad \Leftrightarrow \quad |\Phi_{i_1} \cdots \Phi_{i_L}\rangle, \quad (2.1)$$

where Φ_i on the RHS are 6-dimensional elementary vectors with components $(\Phi_i)_j = \delta_{i,j}$. The one-loop anomalous dimensions of these operators are described by the integrable $SO(6)$ quantum spin-chain Hamiltonian [3]

$$\Gamma = \frac{\lambda}{8\pi^2} H, \quad H = \sum_{l=1}^L \left(1 - \mathcal{P}_{l,l+1} + \frac{1}{2} K_{l,l+1} \right), \quad (2.2)$$

where $\lambda = g_{YM}^2 N$ is the 't Hooft coupling, $\mathcal{P}$ is the permutation operator,

$$\mathcal{P} \Phi_i \otimes \Phi_j = \Phi_j \otimes \Phi_i, \quad (2.3)$$

and the projector K acts as

$$K \Phi_i \otimes \Phi_j = \delta_{ij} \left(\sum_{k=1}^6 \Phi_k \otimes \Phi_k \right). \quad (2.4)$$

It is convenient to define the complex combinations

$$X = \Phi_1 + i\Phi_2, \quad Y = \Phi_3 + i\Phi_4, \quad Z = \Phi_5 + i\Phi_6, \quad (2.5)$$

and to denote the corresponding complex conjugates with a bar, $\bar{X} = \Phi_1 - i\Phi_2$, etc. For $\phi_1, \phi_2 \in \{X, \bar{X}, Y, \bar{Y}, Z, \bar{Z}\}$,

$$\mathcal{P} \phi_1 \otimes \phi_2 = \phi_2 \otimes \phi_1, \quad (2.6)$$

and

$$K \phi_1 \otimes \phi_2 = \begin{cases} 0 & \text{if } \phi_1 \neq \bar{\phi}_2 \\ X \otimes \bar{X} + \bar{X} \otimes X + Y \otimes \bar{Y} + \bar{Y} \otimes Y + Z \otimes \bar{Z} + \bar{Z} \otimes Z & \text{if } \phi_1 = \bar{\phi}_2 \end{cases}. \quad (2.7)$$

2.1 Coordinate Bethe ansatz

We take $|Z^L\rangle$ as the vacuum state, which evidently is an eigenstate of H with zero energy. One-particle excited states (“magnons”) with momentum p are given by

$$|\psi(p)\rangle_\phi = \sum_{x=1}^L e^{ipx} |x\rangle_\phi, \quad (2.8)$$

where

$$|x\rangle_\phi = \left| \overset{1}{\downarrow} Z \cdots Z \overset{x}{\downarrow} \phi Z \cdots Z \overset{L}{\downarrow} Z \right\rangle \quad (2.9)$$

is the state obtained from the vacuum by replacing a single Z at site x with an “impurity” ϕ , which can be either $X, \bar{X}, Y, \bar{Y}$ (but not $\bar{Z}$, which can be regarded as a two-particle bound state). Indeed, one can easily check that (2.8) is an eigenstate of H with eigenvalue $E = \epsilon(p)$, where

$$\epsilon(p) = 4 \sin^2(p/2). \quad (2.10)$$

In order to compute the two-particle S -matrix, we must construct all possible two-particle eigenstates. Let

$$|x_1, x_2\rangle_{\phi_1 \phi_2} = \left| \overset{1}{\downarrow} Z \cdots \overset{x_1}{\downarrow} \phi_1 \cdots \overset{x_2}{\downarrow} \phi_2 \cdots \overset{L}{\downarrow} Z \right\rangle \quad (2.11)$$

denote the state obtained from the vacuum by replacing the Z ’s at sites x_1 and x_2 with impurities ϕ_1 and ϕ_2 , respectively, where $x_1 < x_2$. Following Berenstein and Vázquez [5], we distinguish the following three cases:

$\phi_1 = \phi_2$:

The case of two particles of the same type (i.e., $\phi_1 = \phi_2 \equiv \phi \in \{X, \bar{X}, Y, \bar{Y}\}$) is equivalent to the well-known case originally considered by Bethe in his seminal investigation of the Heisenberg model. (See, e.g., the review by Plefka in [6].) The two-particle eigenstates are given by

$$|\psi\rangle = \sum_{x_1 < x_2} f(x_1, x_2) |x_1, x_2\rangle_{\phi\phi} \quad (2.12)$$

where

$$f(x_1, x_2) = e^{i(p_1 x_1 + p_2 x_2)} + S(p_2, p_1) e^{i(p_2 x_1 + p_1 x_2)}. \quad (2.13)$$

Indeed, these states satisfy

$$H|\psi\rangle = E|\psi\rangle \quad (2.14)$$

with

$$E = \epsilon(p_1) + \epsilon(p_2), \quad (2.15)$$

where $\epsilon(p)$ is given by (2.10). It also follows from (2.14) that the S -matrix for $\phi - \phi$ scattering is given by

$$S(p_2, p_1) = \frac{u_2 - u_1 + i}{u_2 - u_1 - i}, \quad (2.16)$$

where $u_j = u(p_j)$ and

$$u(p) = \frac{1}{2} \cot(p/2). \quad (2.17)$$

$\phi_1 \neq \bar{\phi}_2$:

If the two particles are not of the same type, but $\phi_1 \neq \bar{\phi}_2$, then the two-particle eigenstates are of the form

$$|\psi\rangle = \sum_{x_1 < x_2} \{f_{\phi_1\phi_2}(x_1, x_2) |x_1, x_2\rangle_{\phi_1\phi_2} + f_{\phi_2\phi_1}(x_1, x_2) |x_1, x_2\rangle_{\phi_2\phi_1}\}, \quad (2.18)$$

where

$$f_{\phi_i\phi_j}(x_1, x_2) = A_{\phi_i\phi_j}(12) e^{i(p_1x_1 + p_2x_2)} + A_{\phi_i\phi_j}(21) e^{i(p_2x_1 + p_1x_2)}. \quad (2.19)$$

One finds [5]

$$\begin{pmatrix} A_{\phi_1\phi_2}(21) \\ A_{\phi_2\phi_1}(21) \end{pmatrix} = \begin{pmatrix} R(p_2, p_1) & T(p_2, p_1) \\ T(p_2, p_1) & R(p_2, p_1) \end{pmatrix} \begin{pmatrix} A_{\phi_1\phi_2}(12) \\ A_{\phi_2\phi_1}(12) \end{pmatrix}, \quad (2.20)$$

where the transmission and reflection amplitudes are given by

$$T(p_2, p_1) = \frac{u_2 - u_1}{u_2 - u_1 - i}, \quad R(p_2, p_1) = \frac{i}{u_2 - u_1 - i}, \quad (2.21)$$

respectively.

$\phi_1 = \bar{\phi}_2$:

In the case $\phi_1 = \bar{\phi}_2 \in \{X, \bar{X}, Y, \bar{Y}\}$, the two-particle eigenstates are given by

$$\begin{aligned} |\psi\rangle &= \sum_{x_1 < x_2} \sum_{\phi=X,Y} \{f_{\phi\bar{\phi}}(x_1, x_2) |x_1, x_2\rangle_{\phi\bar{\phi}} + f_{\bar{\phi}\phi}(x_1, x_2) |x_1, x_2\rangle_{\bar{\phi}\phi}\} \\ &\quad + \sum_{x_1} f_{\bar{Z}}(x_1) |x_1\rangle_{\bar{Z}}, \end{aligned} \quad (2.22)$$

where $f_{\phi_i \phi_j}(x_1, x_2)$ are again given by (2.19), and

$$f_{\bar{Z}}(x_1) = A_{\bar{Z}} e^{i(p_1 + p_2)x_1}. \quad (2.23)$$

One finds [5]

$$\begin{pmatrix} A_{X\bar{X}}(21) \\ A_{\bar{X}X}(21) \\ A_{Y\bar{Y}}(21) \\ A_{\bar{Y}Y}(21) \end{pmatrix} = \begin{pmatrix} R(p_2, p_1) & T(p_2, p_1) & S(p_2, p_1) & S(p_2, p_1) \\ T(p_2, p_1) & R(p_2, p_1) & S(p_2, p_1) & S(p_2, p_1) \\ S(p_2, p_1) & S(p_2, p_1) & R(p_2, p_1) & T(p_2, p_1) \\ S(p_2, p_1) & S(p_2, p_1) & T(p_2, p_1) & R(p_2, p_1) \end{pmatrix} \begin{pmatrix} A_{X\bar{X}}(12) \\ A_{\bar{X}X}(12) \\ A_{Y\bar{Y}}(12) \\ A_{\bar{Y}Y}(12) \end{pmatrix}, \quad (2.24)$$

where

$$\begin{aligned} T(p_2, p_1) &= \frac{(u_2 - u_1)^2}{(u_2 - u_1 - i)(u_2 - u_1 + i)}, \\ R(p_2, p_1) &= \frac{-1}{(u_2 - u_1 - i)(u_2 - u_1 + i)}, \\ S(p_2, p_1) &= \frac{-i(u_2 - u_1)}{(u_2 - u_1 - i)(u_2 - u_1 + i)}. \end{aligned} \quad (2.25)$$

2.2 Comparison with the all-loop S -matrix

We now wish to compare the above scattering amplitudes with the weak-coupling limit of the all-loop $SU(2|2) \otimes SU(2|2)$ S -matrix [8]-[14]. This check has not (to our knowledge) been presented elsewhere, and will serve as a useful guide for the $\mathcal{N} = 6$ CS case. It is convenient to express the latter in terms of two mutually commuting sets of Zamolodchikov-Faddeev operators $A_i^\dagger(p)$, $\tilde{A}_i^\dagger(p)$ ($i = 1, \dots, 4$),

$$\begin{aligned} A_i^\dagger(p_1) A_j^\dagger(p_2) &= \sum_{i', j'} S_0(p_1, p_2) \hat{S}_{ij}^{i'j'}(p_1, p_2) A_{j'}^\dagger(p_2) A_{i'}^\dagger(p_1), \\ \tilde{A}_i^\dagger(p_1) \tilde{A}_j^\dagger(p_2) &= \sum_{i', j'} S_0(p_1, p_2) \hat{S}_{ij}^{i'j'}(p_1, p_2) \tilde{A}_{j'}^\dagger(p_2) \tilde{A}_{i'}^\dagger(p_1), \\ A_i^\dagger(p_1) \tilde{A}_j^\dagger(p_2) &= \tilde{A}_j^\dagger(p_2) A_i^\dagger(p_1). \end{aligned} \quad (2.26)$$

We identify the scalar one-particle states as follows,

$$\begin{aligned} X(p) &= A_1^\dagger(p) \tilde{A}_2^\dagger(p), & \bar{X}(p) &= A_2^\dagger(p) \tilde{A}_1^\dagger(p), \\ Y(p) &= A_2^\dagger(p) \tilde{A}_1^\dagger(p), & \bar{Y}(p) &= A_1^\dagger(p) \tilde{A}_2^\dagger(p). \end{aligned} \quad (2.27)$$

The only non-vanishing amplitudes in the scalar sector are

$$\hat{S}_{aa}^{aa}(p_1, p_2) = A, \quad \hat{S}_{ab}^{ab}(p_1, p_2) = \frac{1}{2}(A - B), \quad \hat{S}_{ab}^{ba}(p_1, p_2) = \frac{1}{2}(A + B), \quad (2.28)$$

where $a, b \in \{1, 2\}$ with $a \neq b$. Here

$$\begin{aligned} A &= \frac{x_2^- - x_1^+}{x_2^+ - x_1^-}, \\ B &= - \left[\frac{x_2^- - x_1^+}{x_2^+ - x_1^-} + 2 \frac{(x_1^- - x_1^+)(x_2^- - x_2^+)(x_2^- + x_1^+)}{(x_1^- - x_2^+)(x_1^- x_2^- - x_1^+ x_2^+)} \right], \end{aligned} \quad (2.29)$$

where $x_i^\pm = x(p_i)^\pm$ with

$$\frac{x^+}{x^-} = e^{ip}, \quad x^+ + \frac{1}{x^+} - x^- - \frac{1}{x^-} = \frac{i}{g}, \quad (2.30)$$

and $g = \sqrt{\lambda}/(4\pi)$. Moreover, the scalar factor is given by

$$S_0(p_1, p_2)^2 = \frac{x_1^- - x_2^+}{x_1^+ - x_2^-} \frac{1 - \frac{1}{x_1^+ x_2^-}}{1 - \frac{1}{x_1^- x_2^+}} \sigma(p_1, p_2)^2, \quad (2.31)$$

where $\sigma(p_1, p_2)$ is the BES dressing factor [12, 14]. In the weak-coupling ($g \rightarrow 0$) limit,

$$x^\pm \rightarrow \frac{1}{g} \left(u \pm \frac{i}{2} \right). \quad (2.32)$$

Therefore

$$A \rightarrow \frac{u_1 - u_2 + i}{u_1 - u_2 - i}, \quad B \rightarrow -1, \quad (2.33)$$

and

$$S_0^2 \rightarrow \frac{u_1 - u_2 - i}{u_1 - u_2 + i}, \quad (2.34)$$

since $\sigma(p_1, p_2) \rightarrow 1$.

For two particles of the same type, the scattering amplitude is evidently given by

$$S(p_1, p_2) \equiv \left(S_0(p_1, p_2) \hat{S}_{aa}^{aa}(p_1, p_2) \right)^2 = S_0^2 A^2 \rightarrow \frac{u_1 - u_2 + i}{u_1 - u_2 - i}, \quad (2.35)$$

in agreement with (2.16).

We now consider the case $\phi_1 \neq \bar{\phi}_2$, e.g.,

$$X(p_1) Y(p_2) = T(p_1, p_2) Y(p_2) X(p_1) + R(p_1, p_2) X(p_2) Y(p_1). \quad (2.36)$$

It follows from (2.26)-(2.28) and (2.33), (2.34) that

$$\begin{aligned} T(p_1, p_2) &= \frac{1}{2} S_0^2 A (A - B) \rightarrow \frac{u_1 - u_2}{u_1 - u_2 - i}, \\ R(p_1, p_2) &= \frac{1}{2} S_0^2 A (A + B) \rightarrow \frac{i}{u_1 - u_2 - i}, \end{aligned} \quad (2.37)$$

in agreement with (2.21).

Finally, we consider the case $\phi_1 = \bar{\phi}_2$, e.g.,

$$\begin{aligned} X(p_1) \bar{X}(p_2) &= T(p_1, p_2) \bar{X}(p_2) X(p_1) + R(p_1, p_2) X(p_2) \bar{X}(p_1) \\ &+ S(p_1, p_2) Y(p_2) \bar{Y}(p_1) + S(p_1, p_2) \bar{Y}(p_2) Y(p_1). \end{aligned} \quad (2.38)$$

It follows from (2.26)-(2.28) and (2.33), (2.34) that

$$\begin{aligned} T(p_1, p_2) &= \frac{1}{4} S_0^2 (A - B)^2 \rightarrow \frac{(u_1 - u_2)^2}{(u_1 - u_2 - i)(u_1 - u_2 + i)}, \\ R(p_1, p_2) &= \frac{1}{4} S_0^2 (A + B)^2 \rightarrow \frac{-1}{(u_1 - u_2 - i)(u_1 - u_2 + i)}, \\ S(p_1, p_2) &= \frac{1}{4} S_0^2 (A - B)(A + B) \rightarrow \frac{i(u_1 - u_2)}{(u_1 - u_2 - i)(u_1 - u_2 + i)}, \end{aligned} \quad (2.39)$$

in agreement with (2.25).¹

In short, the all-loop AdS_5/CFT_4 S -matrix correctly reproduces the $\mathcal{N} = 4$ YM one-loop scalar-sector scattering amplitudes, as expected. In the next section, we perform a similar check of the AdS_4/CFT_3 S -matrix.

3 Two-loop S -matrix in the scalar sector of $\mathcal{N} = 6$ CS

The $\mathcal{N} = 6$ CS theory [18] has a pair of scalar fields $A_i(x)$ ($i = 1, 2$) in the bifundamental representation $(\mathbf{N}, \bar{\mathbf{N}})$ of the $SU(N) \times SU(N)$ gauge group, and another pair of scalar fields $B_i(x)$ ($i = 1, 2$) in the conjugate representation $(\bar{\mathbf{N}}, \mathbf{N})$. These fields can be grouped into $SU(4)$ multiplets $Y^A(x)$,

$$Y^A = (A_1, A_2, B_1^\dagger, B_2^\dagger), \quad Y_A^\dagger = (A_1^\dagger, A_2^\dagger, B_1, B_2). \quad (3.1)$$

Following [19], we associate single-trace gauge-invariant scalar operators with states of an alternating $SU(4)$ quantum spin chain with $2L$ sites,

$$\text{tr } Y^{A_1}(x) Y_{B_1}^\dagger(x) \cdots Y^{A_L}(x) Y_{B_L}^\dagger(x) \Leftrightarrow |Y^{A_1} Y_{B_1}^\dagger \cdots Y^{A_L} Y_{B_L}^\dagger\rangle, \quad (3.2)$$

where Y^A on the RHS are 4-dimensional elementary vectors with components $(Y^A)_j = \delta_{A,j}$. The two-loop anomalous dimensions of these operators are described by the integrable

¹There is a sign discrepancy in $S(p_1, p_2)$ which perhaps can be reconciled by a gauge transformation in (2.38), e.g., $Y \rightarrow -Y$ while leaving others unchanged.

alternating $SU(4)$ quantum spin-chain Hamiltonian [19]

$$\Gamma = \lambda^2 H, \quad H = \sum_{l=1}^{2L} \left(1 - \mathcal{P}_{l,l+2} + \frac{1}{2} \{K_{l,l+1}, \mathcal{P}_{l,l+2}\} \right), \quad (3.3)$$

where $\lambda = N/k$ is the 't Hooft coupling², $\mathcal{P}$ is the permutation operator, and the projector K acts as

$$K Y^A \otimes Y_B^\dagger = \delta_B^A \sum_{C=1}^4 Y^C \otimes Y_C^\dagger, \quad K Y_B^\dagger \otimes Y^A = \delta_B^A \sum_{C=1}^4 Y_C^\dagger \otimes Y^C. \quad (3.4)$$

That is,

$$\begin{aligned} K A_i \otimes A_j^\dagger &= K B_i^\dagger \otimes B_j = \delta_{ij} \sum_{k=1}^2 \left(A_k \otimes A_k^\dagger + B_k^\dagger \otimes B_k \right), \\ K A_i^\dagger \otimes A_j &= K B_i \otimes B_j^\dagger = \delta_{ij} \sum_{k=1}^2 \left(A_k^\dagger \otimes A_k + B_k \otimes B_k^\dagger \right), \\ K A_i \otimes B_j &= K B_i \otimes A_j = K A_i^\dagger \otimes B_j^\dagger = K B_i^\dagger \otimes A_j^\dagger = 0. \end{aligned} \quad (3.5)$$

3.1 Coordinate Bethe ansatz

Following [25, 26], we take the state with L pairs of $(A_1 B_1)$, i.e.,

$$|(A_1 B_1)^L\rangle \quad (3.6)$$

as the vacuum state, which evidently is an eigenstate of H with zero energy. It is convenient to label the $(A_1 B_1)$ pairs by $x \in \{1, \dots, L\}$. There are two types of one-particle excited states with momentum p , called “ $\mathcal{A}$ -particles” and “ $\mathcal{B}$ -particles.” The former are given by

$$|\psi(p)\rangle_\phi^{\mathcal{A}} = \sum_{x=1}^L e^{ipx} |x\rangle_\phi^{\mathcal{A}}, \quad (3.7)$$

where

$$|x\rangle_\phi^{\mathcal{A}} = | (A_1 \overset{1}{\downarrow} B_1) \cdots (\phi \overset{x}{\downarrow} B_1) \cdots (A_1 \overset{L}{\downarrow} B_1) \rangle \quad (3.8)$$

is the state obtained from the vacuum by replacing the A_1 from pair x with an “impurity” ϕ , which can be either A_2 or $B_2^\dagger$ (but not $B_1^\dagger$, which can be regarded as a two-particle bound state). Similarly, the “ $\mathcal{B}$ -particles” are given by

$$|\psi(p)\rangle_\phi^{\mathcal{B}} = \sum_{x=1}^L e^{ipx} |x\rangle_\phi^{\mathcal{B}}, \quad (3.9)$$

²The action has two $SU(N)$ Chern-Simons terms with integer levels k and $-k$, respectively.

where

$$|x\rangle_\phi^{\mathcal{B}} = | (A_1 \overset{1}{\downarrow} B_1) \cdots (A_1 \overset{x}{\downarrow} \phi) \cdots (A_1 \overset{L}{\downarrow} B_1) \rangle \quad (3.10)$$

is the state obtained from the vacuum by replacing the B_1 from pair x with an “impurity” ϕ , which can be either $A_2^\dagger$ or B_2 (but not $A_1^\dagger$, which can be regarded as a two-particle bound state). Indeed, both (3.7) and (3.9) are eigenstates of H with eigenvalue $E = \epsilon(p)$, where $\epsilon(p)$ is given by (2.10).

In order to compute the two-particle S -matrix, we must construct all possible two-particle eigenstates.

3.1.1 $\mathcal{A} - \mathcal{A}$ scattering

Let

$$|x_1, x_2\rangle_{\phi_1 \phi_2}^{\mathcal{A}\mathcal{A}} = | (A_1 \overset{1}{\downarrow} B_1) \cdots (\phi_1 \overset{x_1}{\downarrow} B_1) \cdots (\phi_2 \overset{x_2}{\downarrow} B_1) \cdots (A_1 \overset{L}{\downarrow} B_1) \rangle \quad (3.11)$$

denote the state obtained from the vacuum by replacing the A_1 ’s from pairs x_1 and x_2 with impurities ϕ_1 and ϕ_2 , respectively, where $x_1 < x_2$ and $\phi_i \in \{A_2, B_2^\dagger\}$. We distinguish two cases:

$\phi_1 = \phi_2$:

The case of two $\mathcal{A}$ -particles of the same type (i.e., $\phi_1 = \phi_2 \equiv \phi \in \{A_2, B_2^\dagger\}$) is again the same as in the Heisenberg model. The two-particle eigenstates are given by

$$|\psi\rangle = \sum_{x_1 < x_2} f(x_1, x_2) |x_1, x_2\rangle_{\phi\phi}^{\mathcal{A}\mathcal{A}} \quad (3.12)$$

where $f(x_1, x_2)$ is given by (2.13). These states have energy (2.15), and the S -matrix is again given by (2.16),

$$S(p_2, p_1) = \frac{u_2 - u_1 + i}{u_2 - u_1 - i}. \quad (3.13)$$

$\phi_1 \neq \phi_2$:

If the two $\mathcal{A}$ -particles are not of the same type (e.g., $\phi_1 = A_2, \phi_2 = B_2^\dagger$), then the two-particle eigenstates are of the form

$$|\psi\rangle = \sum_{x_1 < x_2} \{ f_{\phi_1 \phi_2}(x_1, x_2) |x_1, x_2\rangle_{\phi_1 \phi_2}^{\mathcal{A}\mathcal{A}} + f_{\phi_2 \phi_1}(x_1, x_2) |x_1, x_2\rangle_{\phi_2 \phi_1}^{\mathcal{A}\mathcal{A}} \}, \quad (3.14)$$

where $f_{\phi_i \phi_j}(x_1, x_2)$ is again given by (2.19). Since K on these states is zero, the S -matrix is again given by (2.21),

$$T(p_2, p_1) = \frac{u_2 - u_1}{u_2 - u_1 - i}, \quad R(p_2, p_1) = \frac{i}{u_2 - u_1 - i}. \quad (3.15)$$

3.1.2 $\mathcal{B} - \mathcal{B}$ scattering

Let

$$|x_1, x_2\rangle_{\phi_1\phi_2}^{\mathcal{B}\mathcal{B}} = | (A_1 \overset{1}{\downarrow} B_1) \cdots (A_1 \overset{x_1}{\downarrow} \phi_1) \cdots (A_1 \overset{x_2}{\downarrow} \phi_2) \cdots (A_1 \overset{L}{\downarrow} B_1) \rangle \quad (3.16)$$

denote the state obtained from the vacuum by replacing the B_1 's from pairs x_1 and x_2 with impurities ϕ_1 and ϕ_2 , respectively, where $x_1 < x_2$ and $\phi_i \in \{A_2^\dagger, B_2\}$. The eigenstates with two $\mathcal{B}$ -particle are given by expressions similar to those with two $\mathcal{A}$ -particles (namely, (3.12) and (3.14) with $|x_1, x_2\rangle_{\phi_i\phi_j}^{\mathcal{A}\mathcal{A}} \leftrightarrow |x_1, x_2\rangle_{\phi_i\phi_j}^{\mathcal{B}\mathcal{B}}$), and we obtain the same results (3.13), (3.15) for the scattering amplitudes.

3.1.3 $\mathcal{A} - \mathcal{B}$ scattering

In order to analyze $\mathcal{A} - \mathcal{B}$ scattering, we define the states

$$\begin{aligned} |x_1, x_2\rangle_{\phi_1\phi_2}^{\mathcal{A}\mathcal{B}} &= | (A_1 \overset{1}{\downarrow} B_1) \cdots (\phi_1 \overset{x_1}{\downarrow} B_1) \cdots (A_1 \overset{x_2}{\downarrow} \phi_2) \cdots (A_1 \overset{L}{\downarrow} B_1) \rangle, \\ |x_1, x_2\rangle_{\phi_2\phi_1}^{\mathcal{A}\mathcal{B}} &= | (A_1 \overset{1}{\downarrow} B_1) \cdots (A_1 \overset{x_1}{\downarrow} \phi_2) \cdots (\phi_1 \overset{x_2}{\downarrow} B_1) \cdots (A_1 \overset{L}{\downarrow} B_1) \rangle, \end{aligned} \quad (3.17)$$

where $x_1 < x_2$ and $\phi_1 \in \{A_2, B_2^\dagger\}$, $\phi_2 \in \{A_2^\dagger, B_2\}$. We distinguish two cases:

$\phi_1 \neq \phi_2^\dagger$:

If $\phi_1 \neq \phi_2^\dagger$ (e.g., $\phi_1 = A_2, \phi_2 = B_2$), then K on the states (3.17) is zero. As noted in [19], we are left with two decoupled $SU(2)$ chains on the even and odd sites. Hence, there is trivial scattering between $\mathcal{A}$ and $\mathcal{B}$ particles.

$\phi_1 = \phi_2^\dagger$:

If $\phi_1 = \phi_2^\dagger$ (e.g., $\phi_1 = A_2, \phi_2 = A_2^\dagger$), then the eigenstates are given by

$$\begin{aligned} |\psi\rangle &= \sum_{x_1 < x_2} \sum_{\phi=A_2, B_2^\dagger} \left\{ f_{\phi\phi^\dagger}(x_1, x_2) |x_1, x_2\rangle_{\phi\phi^\dagger}^{\mathcal{A}\mathcal{B}} + f_{\phi^\dagger\phi}(x_1, x_2) |x_1, x_2\rangle_{\phi^\dagger\phi}^{\mathcal{A}\mathcal{B}} \right\} \\ &+ \sum_{x_1} \sum_{k=1}^2 \left\{ f_{A_k A_k^\dagger}(x_1) |x_1\rangle_{A_k A_k^\dagger} + f_{B_k^\dagger B_k}(x_1) |x_1\rangle_{B_k^\dagger B_k} \right\}, \end{aligned} \quad (3.18)$$

where

$$|x\rangle_{\phi_i\phi_j} = | (A_1 \overset{1}{\downarrow} B_1) \cdots (\phi_i \overset{x}{\downarrow} \phi_j) \cdots (A_1 \overset{L}{\downarrow} B_1) \rangle \quad (3.19)$$

is the state obtained from the vacuum by replacing the $(A_1 B_1)$ pair at x with $(\phi_i \phi_j)$. We assume that $f_{\phi_i \phi_j}(x_1, x_2)$ are again given by (2.19), and

$$f_{\phi_i \phi_j}(x_1) = A_{\phi_i \phi_j} e^{i(p_1 + p_2)x_1}. \quad (3.20)$$

After a lengthy computation (see the Appendix for further details), we find

$$\begin{pmatrix} A_{A_2 A_2^\dagger}(21) \\ A_{A_2^\dagger A_2}(21) \\ A_{B_2^\dagger B_2}(21) \\ A_{B_2 B_2^\dagger}(21) \end{pmatrix} = \begin{pmatrix} 0 & T(p_2, p_1) & 0 & S(p_2, p_1) \\ T(p_2, p_1) & 0 & S(p_2, p_1) & 0 \\ 0 & S(p_2, p_1) & 0 & T(p_2, p_1) \\ S(p_2, p_1) & 0 & T(p_2, p_1) & 0 \end{pmatrix} \begin{pmatrix} A_{A_2 A_2^\dagger}(12) \\ A_{A_2^\dagger A_2}(12) \\ A_{B_2^\dagger B_2}(12) \\ A_{B_2 B_2^\dagger}(12) \end{pmatrix} \quad (3.21)$$

where

$$T(p_2, p_1) = \frac{u_1 - u_2}{u_1 - u_2 - i}, \quad S(p_2, p_1) = \frac{i}{u_1 - u_2 - i}. \quad (3.22)$$

Note that the scattering is reflectionless.

Similar results can be obtained for $\mathcal{B} - \mathcal{A}$ scattering.

3.2 Comparison with the all-loop S -matrix

We now wish to compare the above scattering amplitudes with the weak-coupling limit of the all-loop $SU(2|2)$ S -matrix [28]. It is convenient to express the latter in terms of two sets of Zamolodchikov-Faddeev operators $\mathcal{A}_i^\dagger(p)$, $\mathcal{B}_i^\dagger(p)$ ($i = 1, \dots, 4$) corresponding to the $\mathcal{A}$, $\mathcal{B}$ particles, respectively,

$$\mathcal{A}_i^\dagger(p_1) \mathcal{A}_j^\dagger(p_2) = \sum_{i', j'} S_0(p_1, p_2) \widehat{S}_{i' j'}^{i j}(p_1, p_2) \mathcal{A}_{j'}^\dagger(p_2) \mathcal{A}_{i'}^\dagger(p_1), \quad (3.23)$$

$$\mathcal{B}_i^\dagger(p_1) \mathcal{B}_j^\dagger(p_2) = \sum_{i', j'} S_0(p_1, p_2) \widehat{S}_{i' j'}^{i j}(p_1, p_2) \mathcal{B}_{j'}^\dagger(p_2) \mathcal{B}_{i'}^\dagger(p_1), \quad (3.24)$$

$$\mathcal{A}_i^\dagger(p_1) \mathcal{B}_j^\dagger(p_2) = \sum_{i', j'} \tilde{S}_0(p_1, p_2) \widehat{S}_{i' j'}^{i j}(p_1, p_2) \mathcal{B}_{j'}^\dagger(p_2) \mathcal{A}_{i'}^\dagger(p_1), \quad (3.25)$$

$$\mathcal{B}_i^\dagger(p_1) \mathcal{A}_j^\dagger(p_2) = \sum_{i', j'} \tilde{S}_0(p_1, p_2) \widehat{S}_{i' j'}^{i j}(p_1, p_2) \mathcal{A}_{j'}^\dagger(p_2) \mathcal{B}_{i'}^\dagger(p_1). \quad (3.26)$$

The absence of $\mathcal{A}_{j'}^\dagger(p_2) \mathcal{B}_{i'}^\dagger(p_1)$ terms on the RHS of (3.25) (and similarly, of $\mathcal{B}_{j'}^\dagger(p_2) \mathcal{A}_{i'}^\dagger(p_1)$ terms on the RHS of (3.26)) means that the scattering is reflectionless.

We identify the scalar one-particle states as follows,

$$\begin{aligned} \mathcal{A}_1^\dagger(p)|0\rangle &= \sum_x e^{ipx} |x\rangle_{A_2}^{\mathcal{A}}, & \mathcal{A}_2^\dagger(p)|0\rangle &= \sum_x e^{ipx} |x\rangle_{B_2^\dagger}^{\mathcal{A}}, \\ \mathcal{B}_1^\dagger(p)|0\rangle &= \sum_x e^{ipx} |x\rangle_{B_2}^{\mathcal{B}}, & \mathcal{B}_2^\dagger(p)|0\rangle &= \sum_x e^{ipx} |x\rangle_{A_2^\dagger}^{\mathcal{B}}. \end{aligned} \quad (3.27)$$

The $SU(2|2)$ S -matrix elements $\widehat{S}_{ij}^{i'j'}(p_1, p_2)$ are the same as before (2.28), (2.29), where $x^\pm$ satisfy (2.30) and [25, 26, 27]

$$g = h(\lambda), \quad (3.28)$$

with $h(\lambda) \sim \lambda$ for small λ , and $h(\lambda) \sim \sqrt{\lambda/2}$ for large λ . The scalar factors are given by (cf. (2.31))

$$S_0(p_1, p_2) = \frac{1 - \frac{1}{x_1^+ x_2^-}}{1 - \frac{1}{x_1^- x_2^+}} \sigma(p_1, p_2), \quad \tilde{S}_0(p_1, p_2) = \frac{x_1^- - x_2^+}{x_1^+ - x_2^-} \sigma(p_1, p_2). \quad (3.29)$$

In the weak-coupling ($g \rightarrow 0$) limit,

$$S_0 \rightarrow 1, \quad \tilde{S}_0 \rightarrow \frac{u_1 - u_2 - i}{u_1 - u_2 + i}. \quad (3.30)$$

3.2.1 $\mathcal{A} - \mathcal{A}$ scattering

For two $\mathcal{A}$ particles of the same type (i.e., both $\mathcal{A}_a$ with $a \in \{1, 2\}$), the scattering amplitude is evidently given by

$$S(p_1, p_2) \equiv S_0(p_1, p_2) \widehat{S}_{aa}^{aa}(p_1, p_2) = S_0 A \rightarrow \frac{u_1 - u_2 + i}{u_1 - u_2 - i}, \quad (3.31)$$

in agreement with (3.13). Although the same expression also appears in the $\mathcal{N} = 4$ YM case (2.35), note that the latter follows from the all-loop S -matrix (2.26) in a rather different way.

For two $\mathcal{A}$ particles of different type (i.e., $\mathcal{A}_a$ and $\mathcal{A}_b$ with $a, b \in \{1, 2\}$ and $a \neq b$), it follows from (3.23), (2.28), (2.33) that

$$\mathcal{A}_a^\dagger(p_1) \mathcal{A}_b^\dagger(p_2) = T(p_1, p_2) \mathcal{A}_b^\dagger(p_2) \mathcal{A}_a^\dagger(p_1) + R(p_1, p_2) \mathcal{A}_a^\dagger(p_2) \mathcal{A}_b^\dagger(p_1), \quad (3.32)$$

where

$$\begin{aligned} T(p_1, p_2) &= \frac{1}{2} S_0 (A - B) \rightarrow \frac{u_1 - u_2}{u_1 - u_2 - i}, \\ R(p_1, p_2) &= \frac{1}{2} S_0 (A + B) \rightarrow \frac{i}{u_1 - u_2 - i}, \end{aligned} \quad (3.33)$$

in agreement with (3.15). Again, the same expressions arise in the $\mathcal{N} = 4$ YM case (2.37) in a different way.

3.2.2 $\mathcal{B} - \mathcal{B}$ scattering

According to (3.24), the $\mathcal{B} - \mathcal{B}$ and $\mathcal{A} - \mathcal{A}$ scattering amplitudes are equal, in agreement with the results from Sec. 3.1.2.

3.2.3 $\mathcal{A} - \mathcal{B}$ scattering

According to (3.25), the $\mathcal{A}_a - \mathcal{B}_a$ scattering amplitude is

$$\tilde{S}_0(p_1, p_2) \hat{S}_{aa}^{aa}(p_1, p_2) = \tilde{S}_0 A \rightarrow 1, \quad (3.34)$$

in agreement with the results from Sec.3.1.3 for the case $\phi_1 \neq \phi_2^\dagger$. Note that the scalar factor $\tilde{S}_0$ (3.29) is essential for obtaining this result.

For $\mathcal{A}_a - \mathcal{B}_b$ scattering (with $a, b \in \{1, 2\}$ and $a \neq b$), it follows from (3.25) that

$$\mathcal{A}_a^\dagger(p_1) \mathcal{B}_b^\dagger(p_2) = T(p_1, p_2) \mathcal{B}_b^\dagger(p_2) \mathcal{A}_a^\dagger(p_1) + S(p_1, p_2) \mathcal{B}_a^\dagger(p_2) \mathcal{A}_b^\dagger(p_1), \quad (3.35)$$

where

$$\begin{aligned} T(p_1, p_2) &= \frac{1}{2} \tilde{S}_0 (A - B) \rightarrow \frac{u_1 - u_2}{u_1 - u_2 + i}, \\ S(p_1, p_2) &= \frac{1}{2} \tilde{S}_0 (A + B) \rightarrow \frac{i}{u_1 - u_2 + i}, \end{aligned} \quad (3.36)$$

which agrees with (3.22).³

4 Discussion

We have found that the all-loop AdS_4/CFT_3 S -matrix (3.23) - (3.26) correctly reproduces the $\mathcal{N} = 6$ CS two-loop scalar-sector scattering amplitudes. The scalar factors (3.29), which differ from the AdS_5/CFT_4 scalar factor (2.31), play a crucial role. In particular, we have confirmed that the scattering of $\mathcal{A}$ and $\mathcal{B}$ particles is reflectionless. This gives greater confidence in the correctness of the all-loop S -matrix, and in the corresponding all-loop BAEs [24].

We have restricted our analysis to the scalar sector of $\mathcal{N} = 6$ CS, since this is the only sector for which an explicit Hamiltonian has been available [19]. Very recently, the Hamiltonian for the full two-loop $OSp(6|4)$ spin chain has been found [31, 32]. Hence, it should now be possible to extend the present analysis to other sectors, and thereby further check the all-loop S -matrix.

It would also be interesting to extend the present analysis beyond two loops. This could provide further information about the important function $h(\lambda)$ (3.28) and the dressing phase in the S -matrix. However, such an analysis must wait until the higher-loop Hamiltonian becomes available.

³There is a sign discrepancy in $S(p_1, p_2)$. However, the sign of $S(p_1, p_2)$ in (3.35) can be changed by a gauge transformation, e.g. by changing $\mathcal{A}_1 \rightarrow -\mathcal{A}_1$ and leaving $\mathcal{A}_2, \mathcal{B}_1, \mathcal{B}_2$ unchanged.

Acknowledgments

C.A. thanks the University of Miami for hospitality during the course of this work. This work was supported in part by KRF-2007-313-C00150 and WCU grant R32-2008-000-10130-0 (C.A.), and by the National Science Foundation under Grants PHY-0244261 and PHY-0554821 (R.N.).

A Details of $\mathcal{A} - \mathcal{B}$ scattering

In order to determine the $\mathcal{A} - \mathcal{B}$ scattering amplitudes, it is necessary to act with the Hamiltonian H (3.3) on the state (3.18). We catalog here the action of H on the various terms:

$$\begin{aligned} H|x_1, x_2\rangle_{\phi_i, \phi_j}^{\mathcal{AB}} &= 4|x_1, x_2\rangle_{\phi_i, \phi_j}^{\mathcal{AB}} - |x_1 - 1, x_2\rangle_{\phi_i, \phi_j}^{\mathcal{AB}} - |x_1 + 1, x_2\rangle_{\phi_i, \phi_j}^{\mathcal{AB}} \\ &\quad - |x_1, x_2 - 1\rangle_{\phi_i, \phi_j}^{\mathcal{AB}} - |x_1, x_2 + 1\rangle_{\phi_i, \phi_j}^{\mathcal{AB}} \quad \text{for } x_1 < x_2 - 1, \end{aligned} \quad (\text{A.1})$$

$$\begin{aligned} H|x_1, x_1 + 1\rangle_{A_2, A_2^\dagger}^{\mathcal{AB}} &= 4|x_1, x_1 + 1\rangle_{A_2, A_2^\dagger}^{\mathcal{AB}} - |x_1\rangle_{A_2 A_2^\dagger} - |x_1 + 1\rangle_{A_2 A_2^\dagger} \\ &\quad - |x_1 - 1, x_1 + 1\rangle_{A_2, A_2^\dagger}^{\mathcal{AB}} - |x_1, x_1 + 2\rangle_{A_2, A_2^\dagger}^{\mathcal{AB}}, \end{aligned} \quad (\text{A.2})$$

$$\begin{aligned} H|x_1, x_1 + 1\rangle_{B_2^\dagger, B_2}^{\mathcal{AB}} &= 4|x_1, x_1 + 1\rangle_{B_2^\dagger, B_2}^{\mathcal{AB}} - |x_1\rangle_{B_2^\dagger B_2} - |x_1 + 1\rangle_{B_2^\dagger B_2} \\ &\quad - |x_1 - 1, x_1 + 1\rangle_{B_2^\dagger, B_2}^{\mathcal{AB}} - |x_1, x_1 + 2\rangle_{B_2^\dagger, B_2}^{\mathcal{AB}}, \end{aligned} \quad (\text{A.3})$$

$$\begin{aligned} H|x_1, x_1 + 1\rangle_{A_2^\dagger, A_2}^{\mathcal{AB}} &= 4|x_1, x_1 + 1\rangle_{A_2^\dagger, A_2}^{\mathcal{AB}} - \frac{1}{2}|x_1\rangle_{A_2^\dagger A_2} - \frac{1}{2}|x_1 + 1\rangle_{A_2 A_2^\dagger} \\ &\quad + \frac{1}{2}|x_1\rangle_{A_1 A_1^\dagger} + \frac{1}{2}|x_1 + 1\rangle_{A_1 A_1^\dagger} + \frac{1}{2}|x_1\rangle_{B_1^\dagger B_1} + \frac{1}{2}|x_1 + 1\rangle_{B_1^\dagger B_1} \\ &\quad + \frac{1}{2}|x_1\rangle_{B_2^\dagger B_2} + \frac{1}{2}|x_1 + 1\rangle_{B_2^\dagger B_2} \\ &\quad - |x_1 - 1, x_1 + 1\rangle_{A_2^\dagger, A_2}^{\mathcal{AB}} - |x_1, x_1 + 2\rangle_{A_2^\dagger, A_2}^{\mathcal{AB}}, \end{aligned} \quad (\text{A.4})$$

$$\begin{aligned} H|x_1, x_1 + 1\rangle_{B_2, B_2^\dagger}^{\mathcal{AB}} &= 4|x_1, x_1 + 1\rangle_{B_2, B_2^\dagger}^{\mathcal{AB}} - \frac{1}{2}|x_1\rangle_{B_2^\dagger B_2} - \frac{1}{2}|x_1 + 1\rangle_{B_2^\dagger B_2} \\ &\quad + \frac{1}{2}|x_1\rangle_{B_1^\dagger B_1} + \frac{1}{2}|x_1 + 1\rangle_{B_1^\dagger B_1} + \frac{1}{2}|x_1\rangle_{A_1 A_1^\dagger} + \frac{1}{2}|x_1 + 1\rangle_{A_1 A_1^\dagger} \\ &\quad + \frac{1}{2}|x_1\rangle_{A_2 A_2^\dagger} + \frac{1}{2}|x_1 + 1\rangle_{A_2 A_2^\dagger} \\ &\quad - |x_1 - 1, x_1 + 1\rangle_{B_2, B_2^\dagger}^{\mathcal{AB}} - |x_1, x_1 + 2\rangle_{B_2, B_2^\dagger}^{\mathcal{AB}}, \end{aligned} \quad (\text{A.5})$$

$$\begin{aligned}
H|x_1\rangle_{A_1A_1^\dagger} &= 3|x_1\rangle_{A_1A_1^\dagger} - \frac{1}{2}|x_1-1\rangle_{A_1A_1^\dagger} - \frac{1}{2}|x_1+1\rangle_{A_1A_1^\dagger} \\
&+ \frac{1}{2}|x_1\rangle_{A_2A_2^\dagger} + \frac{1}{2}|x_1+1\rangle_{A_2A_2^\dagger} + |x_1\rangle_{B_1^\dagger B_1} + |x_1+1\rangle_{B_1^\dagger B_1} \\
&+ \frac{1}{2}|x_1\rangle_{B_2^\dagger B_2} + \frac{1}{2}|x_1+1\rangle_{B_2^\dagger B_2} + \frac{1}{2}|x_1-1, x_1\rangle_{A_2^\dagger, A_2}^{\mathcal{AB}} \\
&+ \frac{1}{2}|x_1, x_1+1\rangle_{A_2^\dagger, A_2}^{\mathcal{AB}} + \frac{1}{2}|x_1-1, x_1\rangle_{B_2, B_2^\dagger}^{\mathcal{AB}} + \frac{1}{2}|x_1, x_1+1\rangle_{B_2, B_2^\dagger}^{\mathcal{AB}}, \quad (\text{A.6})
\end{aligned}$$

$$\begin{aligned}
H|x_1\rangle_{A_2A_2^\dagger} &= 4|x_1\rangle_{A_2A_2^\dagger} + \frac{1}{2}|x_1-1\rangle_{A_1A_1^\dagger} + \frac{1}{2}|x_1\rangle_{A_1A_1^\dagger} \\
&+ \frac{1}{2}|x_1\rangle_{B_1^\dagger B_1} + \frac{1}{2}|x_1+1\rangle_{B_1^\dagger B_1} - |x_1-1, x_1\rangle_{A_2, A_2^\dagger}^{\mathcal{AB}} - |x_1, x_1+1\rangle_{A_2, A_2^\dagger}^{\mathcal{AB}} \\
&- \frac{1}{2}|x_1-1, x_1\rangle_{A_2^\dagger, A_2}^{\mathcal{AB}} - \frac{1}{2}|x_1, x_1+1\rangle_{A_2^\dagger, A_2}^{\mathcal{AB}} \\
&+ \frac{1}{2}|x_1-1, x_1\rangle_{B_2, B_2^\dagger}^{\mathcal{AB}} + \frac{1}{2}|x_1, x_1+1\rangle_{B_2, B_2^\dagger}^{\mathcal{AB}}, \quad (\text{A.7})
\end{aligned}$$

$$\begin{aligned}
H|x_1\rangle_{B_1^\dagger B_1} &= 3|x_1\rangle_{B_1^\dagger B_1} - \frac{1}{2}|x_1-1\rangle_{B_1^\dagger B_1} - \frac{1}{2}|x_1+1\rangle_{B_1^\dagger B_1} \\
&+ \frac{1}{2}|x_1-1\rangle_{B_2^\dagger B_2} + \frac{1}{2}|x_1\rangle_{B_2^\dagger B_2} + |x_1-1\rangle_{A_1A_1^\dagger} + |x_1\rangle_{A_1A_1^\dagger} \\
&+ \frac{1}{2}|x_1-1\rangle_{A_2A_2^\dagger} + \frac{1}{2}|x_1\rangle_{A_2A_2^\dagger} + \frac{1}{2}|x_1-1, x_1\rangle_{A_2^\dagger, A_2}^{\mathcal{AB}} \\
&+ \frac{1}{2}|x_1, x_1+1\rangle_{A_2^\dagger, A_2}^{\mathcal{AB}} + \frac{1}{2}|x_1-1, x_1\rangle_{B_2, B_2^\dagger}^{\mathcal{AB}} + \frac{1}{2}|x_1, x_1+1\rangle_{B_2, B_2^\dagger}^{\mathcal{AB}}, \quad (\text{A.8})
\end{aligned}$$

$$\begin{aligned}
H|x_1\rangle_{B_2^\dagger B_2} &= 4|x_1\rangle_{B_2^\dagger B_2} + \frac{1}{2}|x_1\rangle_{B_1^\dagger B_1} + \frac{1}{2}|x_1+1\rangle_{B_1^\dagger B_1} \\
&+ \frac{1}{2}|x_1-1\rangle_{A_1A_1^\dagger} + \frac{1}{2}|x_1\rangle_{A_1A_1^\dagger} + \frac{1}{2}|x_1-1, x_1\rangle_{A_2^\dagger, A_2}^{\mathcal{AB}} \\
&+ \frac{1}{2}|x_1, x_1+1\rangle_{A_2^\dagger, A_2}^{\mathcal{AB}} - |x_1-1, x_1\rangle_{B_2^\dagger, B_2}^{\mathcal{AB}} - |x_1, x_1+1\rangle_{B_2^\dagger, B_2}^{\mathcal{AB}} \\
&- \frac{1}{2}|x_1-1, x_1\rangle_{B_2, B_2^\dagger}^{\mathcal{AB}} - \frac{1}{2}|x_1, x_1+1\rangle_{B_2, B_2^\dagger}^{\mathcal{AB}}. \quad (\text{A.9})
\end{aligned}$$

The appearance of terms of the form $|x\rangle_{A_kA_k^\dagger}$ and $|x\rangle_{B_k^\dagger B_k}$ ($k = 1, 2$) on the RHS of (A.2)-(A.5) explains the need for such terms in the eigenstate (3.18).

With the help of the above results, the eigenvalue equation

$$H|\psi\rangle = E|\psi\rangle \quad (\text{A.10})$$

with $|\psi\rangle$ and E given by (3.18) and (2.15), respectively, leads to the following equations for the amplitudes:

$$\begin{aligned}
0 &= \left[3 - \frac{1}{2}(e^{i(p_1+p_2)} + e^{-i(p_1+p_2)}) - E \right] A_{A_1 A_1^\dagger} \\
&+ (1 + e^{i(p_1+p_2)}) \left(\frac{1}{2} A_{A_2 A_2^\dagger} + A_{B_1^\dagger B_1} + \frac{1}{2} A_{B_2^\dagger B_2} \right) \\
&+ \frac{1}{2}(e^{ip_2} + e^{-ip_1}) \left[A_{A_2^\dagger A_2}(12) + A_{B_2 B_2^\dagger}(12) \right] \\
&+ \frac{1}{2}(e^{ip_1} + e^{-ip_2}) \left[A_{A_2^\dagger A_2}(21) + A_{B_2 B_2^\dagger}(21) \right], \tag{A.11}
\end{aligned}$$

$$\begin{aligned}
0 &= \frac{1}{2}(1 + e^{-i(p_1+p_2)}) A_{A_1 A_1^\dagger} + (4 - E) A_{A_2 A_2^\dagger} + \frac{1}{2}(1 + e^{i(p_1+p_2)}) A_{B_1^\dagger B_1} \\
&+ (e^{ip_2} + e^{-ip_1}) \left[-A_{A_2 A_2^\dagger}(12) - \frac{1}{2} A_{A_2^\dagger A_2}(12) + \frac{1}{2} A_{B_2 B_2^\dagger}(12) \right] \\
&+ (e^{ip_1} + e^{-ip_2}) \left[-A_{A_2 A_2^\dagger}(21) - \frac{1}{2} A_{A_2^\dagger A_2}(21) + \frac{1}{2} A_{B_2 B_2^\dagger}(21) \right], \tag{A.12}
\end{aligned}$$

$$\begin{aligned}
0 &= (1 + e^{-i(p_1+p_2)}) \left[A_{A_1 A_1^\dagger} + \frac{1}{2} A_{A_2 A_2^\dagger} + \frac{1}{2} A_{B_2^\dagger B_2} \right] \\
&+ \left[3 - \frac{1}{2}(e^{i(p_1+p_2)} + e^{-i(p_1+p_2)}) - E \right] A_{B_1^\dagger B_1} \\
&+ \frac{1}{2}(e^{ip_2} + e^{-ip_1}) \left[A_{A_2^\dagger A_2}(12) + A_{B_2 B_2^\dagger}(12) \right] \\
&+ \frac{1}{2}(e^{ip_1} + e^{-ip_2}) \left[A_{A_2^\dagger A_2}(21) + A_{B_2 B_2^\dagger}(21) \right], \tag{A.13}
\end{aligned}$$

$$\begin{aligned}
0 &= \frac{1}{2}(1 + e^{-i(p_1+p_2)}) A_{A_1 A_1^\dagger} + \frac{1}{2}(1 + e^{i(p_1+p_2)}) A_{B_1^\dagger B_1} + (4 - E) A_{B_2^\dagger B_2} \\
&+ (e^{ip_2} + e^{-ip_1}) \left[-A_{B_2^\dagger B_2}(12) + \frac{1}{2} A_{A_2^\dagger A_2}(12) - \frac{1}{2} A_{B_2 B_2^\dagger}(12) \right] \\
&+ (e^{ip_1} + e^{-ip_2}) \left[-A_{B_2^\dagger B_2}(21) + \frac{1}{2} A_{A_2^\dagger A_2}(21) - \frac{1}{2} A_{B_2 B_2^\dagger}(21) \right], \tag{A.14}
\end{aligned}$$

$$\begin{aligned}
0 &= e^{ip_2}(4 - e^{-ip_1} - e^{ip_2} - E) A_{A_2 A_2^\dagger}(12) + e^{ip_1}(4 - e^{-ip_2} - e^{ip_1} - E) A_{A_2 A_2^\dagger}(21) \\
&- (1 + e^{i(p_1+p_2)}) A_{A_2 A_2^\dagger}, \tag{A.15}
\end{aligned}$$

$$\begin{aligned}
0 &= e^{ip_2}(4 - e^{-ip_1} - e^{ip_2} - E)A_{B_2^\dagger B_2}(12) + e^{ip_1}(4 - e^{-ip_2} - e^{ip_1} - E)A_{B_2^\dagger B_2}(21) \\
&- (1 + e^{i(p_1+p_2)})A_{B_2^\dagger B_2}, \tag{A.16}
\end{aligned}$$

$$\begin{aligned}
0 &= e^{ip_2}(4 - e^{-ip_1} - e^{ip_2} - E)A_{A_2^\dagger A_2}(12) + e^{ip_1}(4 - e^{-ip_2} - e^{ip_1} - E)A_{A_2^\dagger A_2}(21) \\
&+ \frac{1}{2}(1 + e^{i(p_1+p_2)}) \left(A_{A_1 A_1^\dagger} - A_{A_2 A_2^\dagger} + A_{B_1^\dagger B_1} + A_{B_2^\dagger B_2} \right), \tag{A.17}
\end{aligned}$$

$$\begin{aligned}
0 &= e^{ip_2}(4 - e^{-ip_1} - e^{ip_2} - E)A_{B_2 B_2^\dagger}(12) + e^{ip_1}(4 - e^{-ip_2} - e^{ip_1} - E)A_{B_2 B_2^\dagger}(21) \\
&+ \frac{1}{2}(1 + e^{i(p_1+p_2)}) \left(A_{A_1 A_1^\dagger} + A_{A_2 A_2^\dagger} + A_{B_1^\dagger B_1} - A_{B_2^\dagger B_2} \right). \tag{A.18}
\end{aligned}$$

Eliminating $A_{A_k A_k^\dagger}$, $A_{B_k^\dagger B_k}$ ($k = 1, 2$), and then solving for the (21) amplitudes in terms of the (12) amplitudes, we arrive at the results (3.21), (3.22).

References

- [1] A. B. Zamolodchikov and Al. B. Zamolodchikov, “Factorized S matrices in two-dimensions as the exact solutions of certain relativistic quantum field models,” *Ann. Phys.* **120**, 253 (1979).
- [2] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2**, 231 (1998) [arXiv:hep-th/9711200]. • S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” *Phys. Lett.* **B428**, 105 (1998) [arXiv:hep-th/9802109]. • E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2**, 253 (1998) [arXiv:hep-th/9802150].
- [3] J. A. Minahan and K. Zarembo, “The Bethe-Ansatz for $\mathcal{N} = 4$ Super Yang-Mills,” *JHEP* **0303**, 013 (2003) [arXiv:hep-th/0212208].
- [4] N. Beisert and M. Staudacher, “The $\mathcal{N} = 4$ SYM Integrable Super Spin Chain,” *Nucl. Phys.* **B670**, 439 (2003) [arXiv:hep-th/0307042].
- [5] D. Berenstein and S. E. Vázquez, “Integrable open spin chains from giant gravitons,” *JHEP* **0506**, 059 (2005) [arXiv:hep-th/0501078].
- [6] A. A. Tseytlin, “Spinning strings and AdS/CFT duality,” in Ian Kogan Memorial Volume, *From Fields to Strings: Circumnavigating Theoretical Physics*, M. Shifman, A. Vainshtein, and J. Wheeler, eds. (World Scientific, 2004) [arXiv:hep-th/0311139]. • N.

- Beisert, “The dilatation operator of $\mathcal{N} = 4$ super Yang-Mills theory and integrability,” *Phys. Rept.* **405**, 1 (2005) [arXiv:hep-th/0407277]. • K. Zarembo, “Semiclassical Bethe ansatz and AdS/CFT,” *Comptes Rendus Physique* **5**, 1081 (2004) [*Fortsch. Phys.* **53**, 647 (2005)] [arXiv:hep-th/0411191]. • J. Plefka, “Spinning strings and integrable spin chains in the AdS/CFT correspondence,” *Living Rev. Rel.* **8**, 9 (2005) [arXiv:hep-th/0507136]. • J. A. Minahan, “A brief introduction to the Bethe ansatz in $\mathcal{N} = 4$ super-Yang-Mills,” *J. Phys.* **A39**, 12657 (2006). • K. Okamura, “Aspects of Integrability in AdS/CFT Duality,” [arXiv:0803.3999].
- [7] M. Staudacher, “The factorized S -matrix of CFT/AdS,” *JHEP* **0505**, 054 (2005) [arXiv:hep-th/0412188].
- [8] N. Beisert, “The $su(2|2)$ dynamic S -matrix,” *Adv. Theor. Math. Phys.* **12**, 945 (2008) [arXiv:hep-th/0511082]. • N. Beisert, “The Analytic Bethe Ansatz for a Chain with Centrally Extended $su(2|2)$ Symmetry,” *J. Stat. Mech.* **0701**, P017 (2007) [arXiv:nlin/0610017].
- [9] R. A. Janik, “The $AdS_5 \times S^5$ superstring worldsheet S -matrix and crossing symmetry,” *Phys. Rev.* **D73**, 086006 (2006) [arXiv:hep-th/0603038].
- [10] G. Arutyunov and S. Frolov, “On $AdS_5 \times S^5$ string S -matrix,” *Phys. Lett.* **B639**, 378 (2006) [arXiv:hep-th/0604043].
- [11] N. Beisert, R. Hernandez and E. Lopez, “A crossing-symmetric phase for $AdS_5 \times S^5$ strings,” *JHEP* **0611**, 070 (2006) [arXiv:hep-th/0609044].
- [12] N. Beisert, B. Eden and M. Staudacher, “Transcendentality and crossing,” *J. Stat. Mech.* **0701**, P021 (2007) [arXiv:hep-th/0610251].
- [13] G. Arutyunov, S. Frolov and M. Zamaklar, “The Zamolodchikov-Faddeev algebra for $AdS_5 \times S^5$ superstring,” *JHEP* **0704**, 002 (2007) [arXiv:hep-th/0612229].
- [14] N. Dorey, D.M. Hofman and J.M. Maldacena, “On the singularities of the magnon S -matrix,” *Phys. Rev.* **D76**, 025011 (2007) [arXiv:hep-th/0703104].
- [15] M. J. Martins and C.S. Melo, “The Bethe ansatz approach for factorizable centrally extended S -matrices,” *Nucl. Phys.* **B785**, 246 (2007) [arXiv:hep-th/0703086].
- [16] M. de Leeuw, “Coordinate Bethe Ansatz for the String S -Matrix,” *J. Phys.* **A40**, 14413 (2007) [arXiv:0705.2369].
- [17] N. Beisert and M. Staudacher, “Long-range $PSU(2, 2|4)$ Bethe ansaetze for gauge theory and strings,” *Nucl. Phys.* **B727**, 1 (2005) [arXiv:hep-th/0504190].

- [18] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, “ $\mathcal{N} = 6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals,” *JHEP* **0810**, 091 (2008) [arXiv:0806.1218].
- [19] J. A. Minahan and K. Zarembo, “The Bethe ansatz for superconformal Chern-Simons,” *JHEP* **0809**, 040 (2008) [arXiv:0806.3951]
- [20] D. Bak and S. J. Rey, “Integrable Spin Chain in Superconformal Chern-Simons Theory,” *JHEP* **0810**, 053 (2008) [arXiv:0807.2063]
- [21] G. Arutyunov and S. Frolov, “Superstrings on $AdS_4 \times CP^3$ as a Coset Sigma-model,” *JHEP* **0809**, 129 (2008) [arXiv:0806.4940].
- [22] B. J. Stefanski, “Green-Schwarz action for Type IIA strings on $AdS_4 \times CP^3$,” [arXiv:0806.4948].
- [23] N. Gromov and P. Vieira, “The AdS_4/CFT_3 algebraic curve,” [arXiv:0807.0437].
- [24] N. Gromov and P. Vieira, “The all loop AdS_4/CFT_3 Bethe ansatz,” *JHEP* **0901**, 016 (2009) [arXiv:0807.0777].
- [25] T. Nishioka and T. Takayanagi, “On Type IIA Penrose Limit and $\mathcal{N} = 6$ Chern-Simons Theories,” *JHEP* **0808**, 001 (2008) [arXiv:0806.3391].
- [26] D. Gaiotto, S. Giombi and X. Yin, “Spin Chains in $\mathcal{N} = 6$ Superconformal Chern-Simons-Matter Theory,” [arXiv:0806.4589].
- [27] G. Grignani, T. Harmark and M. Orselli, “The $SU(2) \times SU(2)$ sector in the string dual of $\mathcal{N} = 6$ superconformal Chern-Simons theory,” [arXiv:0806.4959].
- [28] C. Ahn and R. I. Nepomechie, “ $\mathcal{N} = 6$ super Chern-Simons theory S -matrix and all-loop Bethe ansatz equations,” *JHEP* **0809**, 010 (2008) [arXiv:0807.1924].
- [29] C. Ahn and R. I. Nepomechie, “An alternative S -matrix for $\mathcal{N} = 6$ Chern-Simons theory ?” [arXiv:0810.1915].
- [30] M. Benna, I. Klebanov, T. Klose and M. Smedback, “Superconformal Chern-Simons Theories and AdS_4/CFT_3 Correspondence,” *JHEP* **0809**, 072 (2008) [arXiv:0806.1519].
 • K. Hanaki and H. Lin, “M2-M5 Systems in $\mathcal{N} = 6$ Chern-Simons Theory,” *JHEP* **0809**, 067 (2008) [arXiv:0807.2074]. • I. Shenderovich, “Giant magnons in AdS_4/CFT_3 : dispersion, quantization and finite-size corrections,” [arXiv:0807.2861]. • C. Ahn, P. Bozhilov and R. C. Rashkov, “Neumann-Rosochatius integrable system for strings on $AdS_4 \times CP^3$,” *JHEP* **0809**, 017 (2008) [arXiv:0807.3134]. • Y. Honma, S. Iso,

Y. Sumitomo, H. Umetsu and S. Zhang, “Generalized Conformal Symmetry and Recovery of $SO(8)$ in Multiple M2 and D2 Branes,” [arXiv:0807.3825]. • T. McLoughlin and R. Roiban, “Spinning strings at one-loop in $AdS_4 \times P^3$,” *JHEP* **0812**, 101 (2008) [arXiv:0807.3965]. • J. Kluson, “D2 to M2 Procedure for D2-Brane DBI Effective Action,” *Nucl. Phys.* **B808**, 260 (2009) [arXiv:0807.4054]. • L. F. Alday, G. Arutyunov and D. Bykov, “Semiclassical Quantization of Spinning Strings in $AdS_4 \times CP^3$,” *JHEP* **0811**, 089 (2008) [arXiv:0807.4400]. • C. Krishnan, “AdS4/CFT3 at One Loop,” *JHEP* **0809**, 092 (2008) [arXiv:0807.4561]. • N. Gromov and V. Mikhaylov, “Comment on the Scaling Function in $AdS_4 \times CP^3$,” [arXiv:0807.4897]. • O. Aharony, O. Bergman and D. L. Jafferis, “Fractional M2-branes,” *JHEP* **0811**, 043 (2008) [arXiv:0807.4924]. • G. Bonelli, A. Tanzini and M. Zabzine, “Topological branes, p-algebras and generalized Nahm equations,” [arXiv:0807.5113]. • M. R. Garousi and A. Ghodsi, “Hydrodynamics of $\mathcal{N} = 6$ Superconformal Chern-Simons Theories at Strong Coupling,” [arXiv:0808.0411]. • R. D’Auria, P. Fre, P. A. Grassi and M. Trigiante, “Superstrings on $AdS_4 \times CP^3$ from Supergravity,” [arXiv:0808.1282]. • D. Fioravanti, P. Grinza and M. Rossi, “The generalised scaling function: a systematic study,” [arXiv:0808.1886]. • D. Berenstein and D. Trancanelli, “Three-dimensional $\mathcal{N} = 6$ SCFT’s and their membrane dynamics,” [arXiv:0808.2503]. • R. C. Rashkov, “A note on the reduction of the $AdS_4 \times CP^3$ string sigma model,” *Phys. Rev.* **D78**, 106012 (2008) [arXiv:0808.3057]. • K. Hosomichi, K. M. Lee, S. Lee, S. Lee, J. Park and P. Yi, “A Nonperturbative Test of M2-Brane Theory,” *JHEP* **0811**, 058 (2008) [arXiv:0809.1771]. • J. Kluson and K. L. Panigrahi, “Defects and Wilson Loops in 3d QFT from D-branes in $AdS(4) \times CP^3$,” [arXiv:0809.3355]. • C.-h. Ahn, “Squashing Gravity Dual of $\mathcal{N} = 6$ Superconformal Chern-Simons Gauge Theory,” [arXiv:0809.3684]. • T. McLoughlin, R. Roiban and A. A. Tseytlin, “Quantum spinning strings in $AdS_4 \times CP^3$: testing the Bethe Ansatz proposal,” *JHEP* **0811**, 069 (2008) [arXiv:0809.4038]. • S. Ryang, “Giant Magnon and Spike Solutions with Two Spins in $AdS_4 \times CP^3$,” *JHEP* **0811**, 084 (2008) [arXiv:0809.5106]. • D. Bombardelli and D. Fioravanti, “Finite-Size Corrections of the $\mathbb{CP}^3$ Giant Magnons: the Lüscher terms,” [arXiv:0810.0704]. • T. Lukowski and O. O. Sax, “Finite size giant magnons in the $SU(2) \times SU(2)$ sector of $AdS_4 \times CP^3$,” *JHEP* **0812**, 073 (2008) [arXiv:0810.1246]. • M. Kreuzer, R. C. Rashkov and M. Schimpf, “Near Flat Space limit of strings on $AdS_4 \times CP^3$,” [arXiv:0810.2008]. • C. Ahn and P. Bozhilov, “Finite-size Effect of the Dyonic Giant Magnons in $\mathcal{N} = 6$ super Chern-Simons Theory,” [arXiv:0810.2079]. • C. Ahn and P. Bozhilov, “M2-brane Perspective on $\mathcal{N} = 6$ Super Chern-Simons Theory at Level k,” *JHEP* **0812**, 049 (2008) [arXiv:0810.2171]. • F. Spill, “Weakly coupled $\mathcal{N} = 4$ Super Yang-Mills and $\mathcal{N} = 6$ Chern-Simons theories from $u(2|2)$ Yangian symmetry,” [arXiv:0810.3897]. • J. Gomis, D. Sorokin and L. Wulff, “The complete $AdS(4) \times CP(3)$

- superspace for the type IIA superstring and D-branes,” [arXiv:0811.1566]. • C. Kristjansen, M. Orselli and K. Zoubos, “Non-planar ABJM Theory and Integrability,” [arXiv:0811.2150]. • P. Sundin, “The $AdS(4) \times CP(3)$ string and its Bethe equations in the near plane wave limit,” [arXiv:0811.2775]. • J. H. Baek, S. Hyun, W. Jang and S. H. Yi, “Membrane Dynamics in Three dimensional $\mathcal{N} = 6$ Supersymmetric Chern-Simons Theory,” [arXiv:0812.1772]. • D. Bak, “Zero Modes for the Boundary Giant Magnons,” [arXiv:0812.2645]. • A. Agarwal, N. Beisert and T. McLoughlin, “Scattering in Mass-Deformed $\mathcal{N} \geq 4$ Chern-Simons Models,” [arXiv:0812.3367].
- [31] B. I. Zwiebel, “Two-loop Integrability of Planar $\mathcal{N} = 6$ Superconformal Chern-Simons Theory,” [arXiv:0901.0411].
- [32] J. A. Minahan, W. Schulgin and K. Zarembo, “Two loop integrability for Chern-Simons theories with $\mathcal{N} = 6$ supersymmetry,” [arXiv:0901.1142].